arXiv:1407.4634v3 [quant-ph] 25 Jul 2014

Comment on *Local Unitary equivalence of multipartite pure qubits states*

Ana M. Martins
Instituto Superior Técnico-IST, Universidade de Lisboa-UL,
1049-001 Lisboa, Portugal

July 29, 2014

Abstract

A Comment on the Letter by B. Kraus *Phys. Rev. Lett.* **104**, 020504 (2010).

Comment on "Local Unitary equivalence of multipartite pure qubits states"

Kraus [1] derives necessary and sufficient conditions for the equivalence of *arbitrary*, n-qbit pure states under Local Unitary (LU) operations. Theorem 1, stating that any two pure n-qubit states, with $\rho_i \neq \frac{\mathbf{1}}{2}$, are LU-equivalent iff their *standard forms* are equal, can be formulated using simply the *trace decomposition* form of a state instead of its *standard form*, as proposed in the letter. By one side, it is simpler to work with the *trace decomposition form* of a state then with its *sorted trace decomposition*, that precedes its *standard form*. By another side we show that this new formulation is equally applicable to pure and mixed states. The proof goes like this.

A pure or mixed state $\rho^{'}$ is LU-equivalent to ρ when there exist Local Unitary operators $U_i, ..., U_n$ such that $\rho^{'} = U_1 \otimes ... \otimes U_n \rho \ U_n^{\dagger} \otimes ... \otimes U_1^{\dagger}$. Then

$$\rho_i^{'} = U_i \rho_i, U_i^{\dagger} \tag{1}$$

where $\rho_i^{'}$ and ρ_i are the reduced states of qubit i associated to the global states $\rho^{'}$ and ρ.

Let V_i be the unitary operator that diagonalizes ρ_i, i.e., $\rho_i = V_i D_i V_i^{\dagger}$ where D_i is a diagonal matrix. Introducing this equality in eq.(1) we conclude that $V_i^{'} = U_i V_i$ is the unitary operator that diagonalizes $\rho_i^{'}$, i.e. $\rho_i^{'} = V_i^{'} D_i V_i^{'\dagger}$, therefore the local unitary U_i is uniquely determined (up to a global phase) by $U_i = V_i^{'} V_i^{\dagger}$. When $\rho_i^{'} = \rho_i$, then $V_i^{'} = V_i$ and $U_i = \mathbf{1}$, is the trivial identity operator. However, there are LU-operators, beside the trivial one $U_i = \mathbf{1}$, named *cyclic* local unitary operators [2, 3], that leave a state, $\rho_i \neq \frac{\mathbf{1}}{2}$, invariant. Such LU operators obey the following commutation relation

$$[\rho_i, U_i] = 0 \tag{2}$$

The set of local unitary operators $U_i \in SU(2)$ obeying the cyclic condition (2) is the stabilizer subgroup of ρ_i. The general form of the non trivial unitary cyclic operators U_j, that comute with the 1-qubit density matrix $\rho_j = \frac{1}{2}(\mathbf{1}_j + \vec{r}(j) \cdot \vec{\sigma}(j))$, where $\vec{r}(j) \neq 0$ is the 3-dimensional Bloch vector, $\vec{\sigma}(j) = (\sigma_1(j), \sigma_2(j), \sigma_3(j)$, and where $\sigma_i(j)(i = 1, 2, 3)$ are the Pauli matrices, was derived in Ref.[4]. There it was shown that $U_j = e^{i\xi_j \vec{n}_j \cdot \vec{\sigma}(j)}$, where $\vec{n}_j = \vec{r}(j)/||\vec{r}(j)||$ and ξ_j is a real continuous parameter such that $0 \leq \xi_j \leq \pi/2$.

The *trace decomposition* ρ_t and $\rho_t^{'}$ of *any* two LU-equivalent states ρ and $\rho^{'}$ is

$$\rho_t^{'} = V_1^{'\dagger} \otimes ... \otimes V_n^{'\dagger} \ \ \rho^{'} \ \ V_n^{'} \otimes ... \otimes V_1^{'} \tag{3}$$

$$\rho_t = V_1^\dagger \otimes ... \otimes V_n^\dagger \;\; \rho \;\; V_n \otimes ... \otimes V_1 \tag{4}$$

From where we easily obtain the relation between the two *trace decomposition* forms

$$\rho_t^{'} = (V_1^{'\dagger} U_1 V_1) \otimes ... \otimes (V_n^{'\dagger} U_n V_n) \;\; \rho_t \;\; (V_n^{'\dagger} U_n V_n)^\dagger \otimes ... \otimes (V_1^{'\dagger} U_1 V_1)^\dagger \tag{5}$$

When $\rho_i^{'} \neq \rho_i$ and $\rho_i \neq \frac{\mathbf{1}}{2}$, for all $i = 1, ..., n$, then, $U_i = V_i^{'} V_i^\dagger \Leftrightarrow V_i^{'\dagger} U_i V_i = \mathbf{1}_i$, and the two *trace decomposition* forms are equal, i.e., $\rho_t^{'} = \rho_t$. Given two arbitrary n-qubit states ρ and $\rho^{'}$ we can check if they are LU-equivalent with the following protocol. After verifying that none of the LU operators is cyclic, we express the density operators ρ and $\rho^{'}$ in the generalized Pauli basis [4], and compute separately ρ_t and $\rho_t^{'}$ using (3) and (4). Then we take the difference $(\rho_t^{'} - \rho_t)$. It will be null iff $\rho^{'}$ and ρ are LU-equivalent.

When there is at least one k such that $\rho_k^{'} = \rho_k \neq \frac{\mathbf{1}}{2}$, then $\bar{U}_k = V_k^\dagger U_k V_k$ is given by

$$\bar{U}_k = \begin{pmatrix} e^{i\omega_k} & 0 \\ 0 & e^{-i\omega_k} \end{pmatrix} \tag{6}$$

where $\omega_k = \xi_k ||\vec{r}(k)||$. Assuming that there is only one U_k obeying condition (2) the *trace decomposition forms*, $\rho_t^{'}$ and ρ_t, are related by

$$\rho_t^{'} = \mathbf{1}_1 \otimes ... \otimes \bar{U}_k \otimes ... \otimes \mathbf{1}_n \;\; \rho_t \;\; \mathbf{1}_1 \otimes ... \otimes \bar{U}_k^\dagger \otimes ... \otimes \mathbf{1}_n \tag{7}$$

To check for the LU-equivalence we compute $\rho_t^{'}$ using eq.(3) with $V_k^{'} = V_k$, and the right side of (7) using (4) and (6). Then we compute the difference,

$$\rho_t^{'} - (\mathbf{1}_1 \otimes ... \otimes \bar{U}_k \otimes ... \otimes \mathbf{1}_n \;\; \rho_t \;\; \mathbf{1}_1 \otimes ... \otimes \bar{U}_k^\dagger \otimes ... \otimes \mathbf{1}_n) \tag{8}$$

It will be null iff $\rho^{'}$ and ρ are LU-equivalent. This procedure is easily generalized when there are more than one local unitary operador obeying condition (2).

In conclusion, there are two advantages of using the *trace decomposition form* and density matrix representation of a state instead of the *standard form* and the wave function representation of a state, as is done in the letter [1]: 1) It can be equally applied to pure and mixed states, 2) It is more straighforward to compute the *trace decomposition form* of a state than its *standard form* as it is proposed in the Letter.

References

[1] B. Kraus, *Phys. Rev. Lett.*, **104**, 020504 (2010).

[2] L. B. Fu., *Europhys. Lett.*, **75**, 1 (2006).

[3] S. Gharibian, H. Kampermann, and D. Bruß, *Quantum Information and Computation*, **9**, 1013 (2009).

[4] A. Martins, arXiv:quant-ph/1307.0948v2 (2014).